arXiv:math/0501505v1 [math.DG] 28 Jan 2005

Ricci Curvature and Singularities of Constant Scalar Curvature Metrics

A. Raouf Chouikha[*]

University of Paris-Nord
LAGA,CNRS UMR 7539
Av. J.B. Clement
Villetaneuse F-93430

---

[*] Chouikha@math.univ-paris13.fr

Abstract

In this work we consider periodic spherically symmetric metrics of constant positive scalar curvature on the n-dimensional cylinder called pseudo-cylindric metrics. These metrics belong to the conformal class $[g_0]$ of the Riemannian product $S^1 \times S^{n-1}$: a circle of length T crossed with the (n-1)-dimensional standard sphere.
Such metrics have a harmonic Riemannian curvature and a non parallel Ricci tensor, except for the cylindric one. Thus, it appears a natural link between them and the Derdzinski metrics which are warped product and classify a family of Riemannian manifolds. These two families actually differ by conformal transformations. Moreover, we are interested in the multiplicity problem of the pseudo-cylindric metrics in $[g_0]$. We also study the existence problem and the number of the Derdzinski metrics.
Furthermore, we prove that the pseudo-cylindric metrics may be expressed in terms of elliptic functions for the dimension $n = 3, 4$ and 6 only, and in terms of automorphic functions for any other dimension. This fact allows us to give new bounds for global estimates.
Finally, we examine the curvature of the asymptotic pseudo-cylindric metrics which are complete singular Yamabe metrics on the standard sphere punctured of k points. We show that any of such (non trivial) metric has a non parallel Ricci tensor.

Mathematical Subjects Classification : 53C21, 53C25, 58G30.

Contents

1 **Introduction and statements of results** 4

2 **Singular Yamabe problem and pseudo-cylindric metrics** 8
2.1 The pseudo-cylindric metrics 8
2.2 Existence of metrics with two singular points 10
2.3 Multiplicity of the pseudo-cylindric metrics 11

3 **Remarks on the multiplicity of the Yamabe solutions** 12
3.1 Yamabe metrics on manifolds 12
3.2 The degenerate case 12
3.3 Stability of the solutions and other multiplicity case 13
3.4 The best constant of Sobolev and Aubin theorem 14

4 **Curvature properties** 16
4.1 The Derdzinski metrics and the harmonic curvature 16
4.2 Curvature property of the pseudo-cylindric metrics 17
4.3 Remark concerning the parallelism of the Ricci tensor 18

5 **On the existence of Derdzinski metrics** 20
5.1 Analysis of the Derdzinski equation 20
5.2 Existence conditions of the Derdzinski metrics 21

6 **Singular points of the pseudo-cylindric metrics: a global estimate** 24
6.1 A bound for the pseudo-cylindric fonctions 24
6.2 General remarks 25

7 **Curvature of the asymptotic pseudo-cylindric metrics** 27
7.1 Characterization of metrics with k singularities 27
7.2 Ricci tensor of the asymptotic pseudo-cylindric metrics 28

8 **Proofs** 30
8.1 Proof of Theorem (2-4) 30
8.2 Proof of Theorem (4-3) 33
8.3 Proof of Theorem (6-1) 35
8.4 Proof of Theorem (7-3) 39

1 Introduction and statements of results

Let us consider a smooth compact Riemannian manifold (M, g) of dimension $n \geq 3$, where g is a smooth Riemannian metric on M. dv denotes the volume form of g, and $R(x)$ its scalar curvature. ${\mathcal{H}_1}^2(M, g)$ is the Sobolev space.
For any non null function $u \in {\mathcal{H}_1}^2(M, g)$, we define the functional

$$J(u) = \frac{4\frac{n-1}{n-2} \int_M |\nabla u|^2 \, dv + \int_M R(x) u^2 dv}{\left(\int_M u^{\frac{2n}{n-2}} dv\right)^{\frac{n-2}{n}}} \ . \tag{1}$$

The critical points of the functional $J(u)$ are solutions of the Euler-Lagrange equation

$$4\frac{n-1}{n-2}\Delta_g u + R(x)u = k|u|^{\frac{4}{n-2}}u \ , \tag{2}$$

where k is a real number.

There is a geometric interest only if we assume the solution u of (2) to be C^∞ positive. In this case, the conformal metric $g' = u^{\frac{4}{n-2}}g$ has constant scalar curvature k if u is a critical point of the functional $J(u)$ and conversely. So, we know by resolution of the Yamabe conjecture for the smooth compact manifolds, the infimum of $J(u)$ is always attained . A such metric with constant scalar curvature belonging to the conformal class $[g]$ is called Yamabe metric.
More precisely, let (S^n, g_0) be the standard sphere, we may assert, [L-P] :

There exists a non zero function u_0, *positive,* C^∞ *on* (M, g) *such that*

$$J(u_0) = Inf_{u \in H_1^2(M)} J(u) = \mu(M, g).$$

Moreover, we have $\mu(M, g) < \mu(S^n, g_0)$, *equality holds only if there exists a conformal diffeomorphism between the manifold* (M, g) *and the standard sphere* (S^n, g_0).

It is also known that the Yamabe metric is unique up to homothety, when (M, g) is such that $\int_M R(x) dV \leq 0$ or when (M, g) is an Einstein Manifold.
For the positive case of the scalar curvature $(k > 0)$, it is possible to get multiple Yamabe solutions for certain conformally flat product Riemannian manifolds.

The multiplicity problem of the Yamabe solutions have been intensively studied during the last few years, in the more general context known as the singular Yamabe problem with isolated singularities. The latter consists to find a metric $g = u^{\frac{4}{n-2}} g_0$ on the domain $S^n - \Lambda_k$, where Λ_k is a finite point-set of (S^n, g_0). This metric being complete and have a constant scalar curvature $R(g)$.

Concerning the special case $k = 2$, R. Schoen (see section 2 of [Sc]) has partially described the conformal class of the product metric $(S^1 \times S^{n-1}, dt^2 + d\xi^2)$, where S^1 is the circle of length T and (S^n, g_0) is the standard sphere.

He proved the existence and the number of their rotationally invariant Yamabe metrics

$$g^j{}_c = u^j{}_c{}^{\frac{4}{n-2}} (dt^2 + d\xi^2) \ ; \quad j = 1, 2, ..., k-1$$

called pseudo-cylindric metrics (sometime also called by different authors Delaunay metrics or Fowler metrics) which depend on T-values. But, his proof is not completed. The completion has not been yet published elsewhere (unless our unpublished joint work with F. Weissler [Ch-W]).
Now, it is well known that the number of these metrics is finite in a conformal class for the product $[dt^2 + d\xi^2]$.
We refer to the papers [Sc], [Ch3] or [Ch-W].
In particular, we get a unique Yamabe metric if

$$T \leq 2\pi \sqrt{\frac{n-1}{R_0}} \ . \tag{3}$$

This metric is $c^{te}(dt^2 + d\xi^2)$.

One of our main results concerns geometric properties of the above metrics. That is any non trivial pseudo-cylindric metric have a Ricci tensor non parallel and so, its Riemannian curvature is harmonic (see Theorem (4-3)below and [Ch3]). A tensor of order two $\tau = (\tau_{ij})$ is said to be nonparallel if the covariant differentiation of τ along (M, g) has a nonzero component.

Observe that all these metrics -sharing the above curvature properties-have been classified by A. Derdzinski. Indeed, these metrics are covered by a warped product metrics : $dt^2 + f^2(t) d\xi^2$ up to an isometry, where $f(t)$ satisfy a certain ODE. Moreover, we show by using a Yau result (Lemma (4-5) below), that any pseudo-cylindric metric may actually be identified with a Derdzinski metric, up to a conformal diffeomorphism. This identification may be written

$$dt^2 + f^2(t) d\xi^2 = F^*(u^j{}_c{}^{\frac{4}{n-2}} (dt^2 + d\xi^2)),$$

where F is a conformal diffeomorphism.
Remark that the metrics originally contructed by A. Derdzinski, permit to provide example of compact Riemannian manifold with harmonic curvature and non-parallel Ricci tensor. We know that these metrics are depending on the S^1- parametrization T (thus, there are obviously not conformally equivalent) .

Nevertheless, the pseudo-cylindric metrics belong to the same conformal class which is parametrized by T. Hence, they are conformally equivalent to the cylindric metric. That is not the same for the Derdzinski metrics. For this reason, from the multiplicity point of view of the Yamabe solutions, the pseudo-cylindric metrics are more interesting. This fact will be clarified below.

This paper is organized as follows :

1 - We achieve in Section 2, the completed proof (not yet published somewhere) concerning the existence and the finite number of the pseudo-cylindric metrics in a conformal class.

2 - In section 3, we consider the link between the multiplicity of Yamabe problem and the best constant of sobolev. In particular, we put right and improve a theorem of T. Aubin.

3 - In Section 4, we examine the non parallelism properties of the Ricci curvature for the pseudo-cylindric metrics.

4 - In the next Section 5 we are interested on the existence of many Derdzinski metrics, satisfying certain conditions. In particular, we show the existence of a finite number (depending on the circle length T) of such metrics.
This fact has not been remarked in the Part D of chapter 16 of A. Besse [Be]. His Theorem 16.38 only proved existence of non trivial Derdzinski metrics without precising the interconnection between these metrics and the circle length.
In particular, we exhibit a constant T_0 such that, if we assume $T \leq T_0$, then there is only one Derdzinski metric with parallel Ricci tensor, i.e. the trivial one (see Theorem (5-3) below).

5 - Furthermore, for the dimension values $n = 3, 4, 6.$ the pseudo-cylindric solution can be expressed in terms of a real periodic elliptic functions. The interest of the latter remark is that an elliptic function has only polar singularities. So, as a corollary (see also our previous work [Ch1]), we deduce the estimates for the corresponding pseudo-cylindric solutions (Theorem (6-1) below):

$$r^{\frac{n-2}{2}} u(r) \leq C, \qquad \textit{and} \qquad r^{\frac{n}{2}} \frac{du(r)}{dr} \leq C' \ .$$

This fact allows us to explicitly determine the bounds which naturally depend on the dimension value n and the period T.
This generalizes a result of Cafarelli-Gidas-Spruck [C-G-S] - According to Pollack (Theorem (4-3) of [P]), the first inequality has been etablished by R. Schoen - .
Moreover, we show for any dimension other than $3, 4$ or 6, the pseudo-cylindric solutions may be expressed in terms of automorphic functions. So, they are invariant under a substitution group.

6 - Finally in section 7, we consider the asymptotic pseudo-cylindric metrics. They are complete Yamabe metrics on $S^n - \Lambda_k$, where $\Lambda_k = \{p_1, p_2, ..., p_k\}$ (here $k > 2$). In particular, we show that for any non trivial metrics, its Ricci tensor is also non parallel (Theorem (7-3) below).

All the proofs are exposed to the end, in Section 8.

2 Singular Yamabe problem and pseudo-cylindric metrics

2.1 The pseudo-cylindric metrics

Let a finite point-set $\Lambda_k = p_1, p_2, ..., p_k$ of the standard sphere S^n. R_0 being its scalar curvature. consider the following

Singular Yamabe problem:
There exists a metric $g = u^{\frac{4}{n-2}} g_0$ *in the conformal class* $[g_0]$ *on the domain* $S^n - \Lambda_k$*, which is complete and has a constant scalar curvature* $R(g)$*.*
In others words, the following differential equation

$$4\frac{n-1}{n-2}\Delta_{g_0} u + R_{g_0} u - R_g u^{\frac{n+2}{n-2}} = 0, \tag{4}$$

$$g = u^{\frac{4}{n-2}} g_0 \quad \text{is complete on} \quad S^n - \Lambda \ , \qquad R_g = \text{constant} > 0,$$

has a solution.

Let $\mathcal{L}_g = \Delta_g + \frac{n-2}{4(n-1)} R_g$ denotes the conformal Laplacian of any Riemannian manifold (M, g). $\mathcal{L}_g$ posseses a special property:

Invariance property of the conformal Laplacian : Let (M, g) *and* (N, h) *two Riemannian manifolds, such that there exists a conformal diffeomorphism* $f : M \rightarrow N$ *(that means if* f *is a diffeomorphism, there is a function* $v : M \rightarrow IR^+$*, such that* $f^* h = v^{\frac{4}{n-2}} g$*).*
Thus, for any function $\varphi : N \rightarrow IR$ *of class* C^2*, we have*

$$\mathcal{L}_h(\varphi) \circ f = v^{\frac{n+2}{n-2}} \mathcal{L}_g(v \varphi \circ f)$$

($\varphi = 1$ *and* $f = Id$ *gives the Yamabe equation).*

Consequently, we may deduce from the above, if v is any positive solution of

$$\mathcal{L}_{g_0} v - \frac{n-2}{4(n-1)} K v^{\frac{n+2}{n-2}} = 0, \tag{5}$$

then K is precisely the scalar curvature of the metric $g = v^{\frac{4}{n-2}} g_0$.
Moreover, condition g complete requires that $lim_{x \rightarrow \Lambda} v(x) = \infty$.

The interest of the problem was underlined by Schoen; he first proved the existence of weak solutions of Yamabe problem on (S^n, g_0), which have a

prescribed singular set Λ_k .
In particular, this latter point allows us to prove the existence of metrics g conformally equivalent to g_0 , which have positive constant scalar curvature and are complete on $S^n - \Lambda_k$. We refer to [M-Pa] for more details about the general problem, and for a simpler version of the Schoen result.

The set Λ_k may be replaced by $F(\Lambda_k) = \Lambda'_k$, for any conformal transformation. In particular, we may suppose that the points p_i sum to zero, considered as vectors in the euclidian space and then Λ_k is contained in an equatorial subsphere $S^k \subset S^n$. By using a reflection argument, one proves there is no complete (positive regular) solution existing for the value $k = 1$. One remarks a similarity between this singular Yamabe problem and the theory of embedded, complete, constant mean curvature surfaces with k ends in $I\!R^3$.
For the case $k = 2$, a family of such surfaces was discovered by C. Delaunay. For this reason, it was called "Delaunay metrics" the analogous family of solutions for the corresponding singular Yamabe problem. Some authors also called them "Fowler metrics" in relation with radial solutions of Emden-Fowler equations.

It is known the moduli space $\mathcal{M}_\Lambda$, which is the set of all smooth positive functions u to the problem

$$4\frac{n-1}{n-2}\Delta_{g_0}u + R(g_0)u - R(g)u^{\frac{n+2}{n-2}} = 0 \tag{6}$$

$$g = u^{\frac{4}{n-2}}g_0 \quad \text{is complete on} \quad S^n - \Lambda \ , \qquad R(g) = \text{constant} > 0.$$

This space is, in general a k -dimensional real analytic manifold, see [M-P-U]. In the general case where $\Lambda = S^p$ (the p-subsphere where $1 \leq p \leq \frac{n-2}{2}$), these solutions belong to an infinite dimensional space. F. Pacard [Pa] has shown the existence of a positive complete solution in the case $p = \frac{n-2}{2}$. Nevertheless, the moduli space is only explicitly determined if $dim\Lambda = 0$, *i.e.* $\Lambda = \Lambda_k$.
In the case $k = 2$, we remark that the space $\mathcal{M}_\Lambda$ may be identified with any other, when Λ contains just two elements. This space, denoted by $\mathcal{M}_2$ which contains all the pseudo-cylindric metrics, may be identified with the open set $\Omega \in I\!R^2$. The frontier of Ω is the homoclinic curve γ_{b_0} , which will be described below.

Since this problem is conformally invariant, we may choose the set $\Lambda_2 = (p, -p)$; $-p$ is the antipodal point of p .

Let $\Pi : S^n - \{pt\} \to IR^n$ the stereographic projection. We know that the converse Π^{-1} defined by

$$\Pi^{-1}(x) = [\frac{2x}{|x|^2+2}, \frac{|x|^2-1}{|x|^2+1}]$$

is a conformal diffeomorphism, and we have $(\Pi^{-1})^\star(g_0) = \frac{4}{(|x|^2+1)^2}dx^2$, where dx^2 is the euclidean metric of IR^n.
Thus, the resolution of (4) is equivalent to the resolution of

$$\Delta u + \frac{n(n-2)}{4}u^{\frac{n+2}{n-2}} = 0 \tag{7}$$

on $IR^n - \tilde{\Lambda}_k$, where $\tilde{\Lambda}_k = \Pi(\Lambda_k)$. Moreover, u have to verify $lim_{x \to \tilde{\Lambda}_k} u(x) = \infty$.

2.2 Existence of metrics with two singular points

Examine now the case $k = 2$. In this case, it will be possible to descrive all the Yamabe metrics. It is known that if u is a solution on $S^n - \{p_1, p_2\}$, then it is invariant with respect to any conformal transformation fixing these points. If these points are antipodal (as we may assume) then u is rotationally invariant. It is then convenient to project the sphere S^n to IR^n, from one of the singular point, p_1, so that p_2 sent to $0 \in IR^n$.
In this case, we have choosen $\tilde{\Lambda}_k = (0, \infty)$.
On the other hand, Cafarelli, Gidas and Spruck [C-G-S] proved that all solutions of (7) on $IR^n - \tilde{\Lambda}$ are radial. We then define the change

$$u(x) = |x|^{\frac{2-n}{2}} v(log\frac{1}{|x|}).$$

In fact, this change corresponds to the conformal diffeomorphism

$$\beta : IR^n - \{0\} \to IR \times S^{n-1}$$

where $\beta(x) = (log\frac{1}{|x|}, \frac{x}{|x|})$.
Consequently, $v : IR \to IR^+$ is a solution of the autonomous ODE

$$\frac{d^2}{dt^2}v - \frac{(n-2)^2}{4}v + \frac{n(n-2)}{4}v^{\frac{n+2}{n-2}} = 0 \tag{8}$$

The analysis of (8) show us, there is only one center say $(\alpha, 0)$ corresponding to the constant solution $\alpha = (\frac{n-2}{n})^{\frac{n-2}{2}}$. All the periodic orbits denoted by

$\gamma_c(t)$, are parametrized by $(v_c(t), \frac{d}{dt}v_c(t))$, c verifying $0 < c < c_0$. They are surrounded by the homoclinic orbit γ_{c_0}. Here $v_{c_0}(t) = (\cosh t)^{\frac{n-2}{2}}$. The period is also depending on c : $T = T(c)$.
The metrics

$$g = {v_c}^{\frac{4}{n-2}}(dt^2 + d\xi^2)$$

are called pseudo-cylindric type metrics. They are defined on the product manifold $(S^1 \times S^{n-1}, dt^2 + d\xi^2)$ a circle of length T crossed with the standard (n-1)-dimensional sphere. They belong to the conformal class of the cylindric metric $[dt^2 + d\xi^2]$.

2.3 Multiplicity of the pseudo-cylindric metrics

R. Schoen [Sc] gave a description of the family of metrics with constant scalar curvature in the conformal class of the product metric $g_0 = dt^2 + d\xi^2$ on $S^1(T) \times S^{n-1}$ the product Riemannian manifold of a circle of length T and the euclidian (n-1)-sphere of radius 1.
Many other authors tried to descrive the conformal class $[g_0]$, in order to complete the Schoen analysis concerning this problem, (see page 134 section 2 of [Sc]), In particular, we refer to [Ch1], and for a more geometrical interpretation we refer to [M-Pa].
The following result resume all the known properties of the pseudo-cylindric metrics, see [Sc], [Ch2], [M-Pa] and [Ch-W]

Theorem (2-4)
Let be $(S^1(T) \times S^{n-1}, g_0 = dt^2 + d\xi^2)$ *the product Riemannian manifold of a circle of length* T *and the euclidian (n-1)-sphere of radius 1. If the value of* T *satisfies the condition*

$$T_{k-1} = \frac{2\pi(k-1)}{\sqrt{n-2}} < T \leq T_k = \frac{2\pi k}{\sqrt{n-2}}, \tag{9}$$

where the integer $k \geq 1$.
Then, there exist, in the conformal class $[g_0]$ *k rotationally invariant pseudo-cylindric metrics*

$${g^j}_c = ({w^j}_c)^{\frac{4}{n-2}}(dt^2 + d\xi^2) \quad , \quad j = 0, 1, 2, .., k-1 \quad \text{with} \quad {g^0}_c = g_0.$$

$k = 1$ *only corresponds to the trivial product metric* $C^{te}(dt^2 + d\xi^2)$.

3 Remarks on the multiplicity of the Yamabe solutions

3.1 Yamabe metrics on manifolds

Recall that for the standard sphere (S^n, g_0) , we only get a trivial result: all Yamabe solutions are minimal for the functional $J(u)$ and their sectional curvature turn to be constant. There actually exists a conformal diffeomorphism between the manifolds $(S^n, u^{\frac{4}{n-2}} g_0)$ and (S^n, g_0) . In fact, all the metrics are obtained by pulling back the canonical one, by a natural family of conformal transformations.

Among these metrics, we have an explicit one parameter family of metrics (of constant scalar curvature R)

$$g_t = (\sqrt{1+t^2} + t\cos\alpha x)^{-2} g_0 \qquad \text{with} \qquad \alpha = \sqrt{\frac{R}{n(n-1)}}. \tag{10}$$

We can verify that all the functions $u_t(x) = (\sqrt{1+t^2} + t\cos\alpha x)^{-\frac{n-2}{2}}$ are solutions of the Yamabe equation (2) for $R \equiv k$. These functions are minimizing for the functional, i.e. $J(u_t) = \mu(S^n, g_0)$.

Moreover, consider a Riemannian manifold (M, g) with constant scalar curvature R which in addition admits a conformal isometric vector field $\mathcal{X}$. Then, $\frac{R}{n-1}$ belongs to the spectre of the Laplace operator Δ_g . By pulling back g by the flow of $\mathcal{X}$ we get a one parameter family of metrics which are conformal to g and have the same scalar curvature R .

3.2 The degenerate case

$T = T_k$ corresponds to the degenerate case which has previously been considered. The product metrics in general are not Morse critical points. Notice that the Morse index of the solution $g = u^{\frac{4}{n-2}} g_0$ is equal to the number of eigenvalues of the Laplace operator Δ_g in $[0, n]$. The assumption that the solution is not degenerate implies that the strict inequality $\lambda_1 > n$ holds. According to Lichnerowicz, we get $\lambda_1 \geq n$, whenever the metric g is Yamabe solution.

More generally, Schoen proves that, if all Yamabe solutions are non degenerate in a conformal class $[g_0]$ of a Riemannian compact manifold (M, g_0) then the functional has only a finite number of Morse critical points. Thus, for a generic conformal class of metrics this hypothesis on g_0 is true, and

$[g_0]$ contains at most a finite number of (non degenerate) Yamabe metrics. Moreover, the set of the Yamabe metrics with finite energy is compact.

3.3 Stability of the solutions and other multiplicity case

(i) Consider the stability problem of the pseudo-cylindric metrics ${g^j}_c = ({u^j}_c)^{\frac{4}{n-2}} g_0$, we want to precise that among the k metrics given by the above theorem, only the solution u_T with fundamental period T is stable. Notice that the limiting period is the fundamental period of the associated linearized equation of

$$\frac{d^2}{dt^2}u + (n-2)u.$$

The other $(k-1)$ solutions which have fundamental period $\frac{T}{j}$ with $j = 2, ..., k-1$, are unstable.

By crossing the point $(T_1, u_{T_1}(t))$, there is an "exchange of stability" between the two curves of solutions. The trivial one becomes unstable after crossing this bifurcation point. Therefore, if we suppose $T > T_1$ the trivial solution $u \equiv \alpha$ does not minimize the Yamabe functional

$$J(u) = \frac{4\frac{n-1}{n-2}\int_M |\nabla u|^2 dv + \int_M R(x)u^2 dv}{\left(\int_M u^{\frac{2n}{n-2}} dv\right)^{\frac{n-2}{n}}} .$$

In fact, the infimum is attained by the solution u_T having the fundamental period T .

If we get $T \leq T_1$, the (unique) solution $u \equiv \alpha$ minimizes the functional. To see that, it is enough to calculate its value.

Indeed, let V denotes the Riemannian volume of $(S^1(T) \times S^{n-1}, g_0)$, we get

$$J(u \equiv \alpha) = (n-1)(n-2)\Big(\int dv\Big)^{\frac{2}{n}}$$

where

$$\alpha = (\frac{n-2}{n})^{\frac{n-2}{4}} \quad and \quad \Big(\int dv\Big) = T vol(S^{n-1}, d\xi^2) .$$

If we suppose

$$T \leq \frac{2\pi}{\sqrt{n-2}} ,$$

obviously, we get

$$J(u \equiv \alpha) < n(n-1)[vol(S^n)]^{\frac{2}{n}} .$$

Furthermore, notice that only the non trivial solution u_T tends to $u_0(t) = (cosht)^{-\frac{n-2}{2}}$, when T becomes ∞ . The minimum of the Yamabe functional μ_T has to tend to $\mu(S^n, g_0) = n(n-1)[vol(S^n)]^{2/n}$, when T tends to ∞ .

(ii) The multiplicity of the Yamabe problem may occur for other locally conformally flat manifolds. Consider the real projective space: among the pseudo-cylindric metrics $g = u^{\frac{4}{n-2}} g_0$, we examine those satisfying the condition $u(t,\xi) = u(-t,-\xi)$; any such solution is independent of ξ . Thus we obtain solutions to the Yamabe problem on the punctured real projective space $RP^n - \{p\}$ by identifying any solution on this space with a solution on $S^n - \{p,-p\}$, invariant under the antipodale reflection.

3.4 The best constant of Sobolev and Aubin theorem

There is another problem related on the multiplicity problem of Yamabe solutions: to determine the best constant of the Sobolev inclusions. Aubin [A1] has shown that for a positive number $A \geq K^2(n,2) = \dfrac{4\omega_n^{-\frac{2}{n}}}{n(n-2)}$ on the compact Riemannian manifold (M,g) *of dimension* $n > 2$, where ω_n denotes the volume of the standard sphere, there exists a constant B such that, for all $u \in H_1^2(M,g)$, we have the following Sobolev inequality

$$\|u\|^2_{L^{\frac{2n}{n-2}}} \leq A\|\nabla u\|^2_{L^2} + B\|u\|^2_{L^2} \ . \tag{11}$$

The existence of the second constant B seems play a role in the Yamabe multiplicity problem and in the Nirenberg problem. Hebey-Vaugon [H-V] have calculated the best little constant B denoted by $C(M,g)$. For this constant precisely, the inequality (11) is verified by any function $u \in H_1^2(M,g)$, on the quotient manifold of the standard sphere S^n/G, where G is a cyclic group of isometries on the sphere.

Moreover, one has estimated the best constant for a conformally flat manifolds $S^1 \times S^{n-1}$ endowed with a Riemannian metric product. More precisely, they find

$$C(S^1(T) \times S^{n-1}) \leq \frac{4\omega_n^{-\frac{2}{n}}}{n(n-2)} T^{-2} + \frac{n-2}{n}\omega_n^{-\frac{2}{n}}$$

where T is the circle radius.

Concerning the first constant A . Notice that A may attain the value $K^2(n,2)$ if the manifold (M,g) has constant sectional curvature.

According to implicit function theorem, the Yamabe equation (4) is not locally invertible on $(S^1(T) \times S^{n-1})$ near the trivial solution $u \equiv \alpha$, and

$$\lambda = (\frac{2\pi}{T})^2 = n-2$$

is an eigenvalue, which yet verify the inequality

$$\lambda < n\Big(\frac{\omega_n}{\int dv}\Big)^{\frac{2}{n}} \ .$$

Obviously, the constant $K^2(n,2)$ is not attained, and there exists $u \in H_1^2(M,g)$ such that

$$\|u\|^2_{L^{\frac{2n}{n-2}}} > K^2(n,2)\Big(\|\nabla u\|^2_{L^2} + \frac{(n-2)^2}{4}\|u\|^2_{L^2}\Big) \ ,$$

since we have $\mu(S^1(T)\times S^{n-1}) < \mu(S^n, g_0) = K^{-2}(n,2)$
Thus, we have proved the following.

Proposition (3-3)
Consider V_n *a Riemannian compact manifold* ($n \geq 3$), *and the equation on* V_n

$$\Delta\phi + k\phi = k(1+f)\phi^{\frac{n+2}{n-2}}.$$

If, for $k = k_0$ *this equation is not locally invertible on* $\phi_0 = 1$, *then* $\lambda = \frac{4k_0}{n-2}$ *is an eigenvalue of the Laplace operator.*
Moreover, if the eigenvalue verify $\lambda \geq n\Big(\frac{\omega_n}{\int dv}\Big)^{\frac{2}{n}}$, *then the Yamabe problem have at least two solutions. Otherwise,* V_n *is conformal to the standard sphere.*

This proposition puts right and completes Theorem 11 of Aubin [A1], in including metrics like as the pseudo-cylindric metrics (with respect to notations of [A1])

4 Curvature properties

4.1 The Derdzinski metrics and the harmonic curvature

Let us consider a compact Riemannian manifold (M, g) , $\mathcal{R}$ is the curvature associated to the metric g , $r = Ric(g)$ is its Ricci tensor, and D is the Riemannian connexion. We denote by $\delta\mathcal{R}$ its formal divergence. This curvature is harmonic if $\delta\mathcal{R} = 0$. According to the second Bianchi identity

$$\delta\mathcal{R} = -d(Ric(g));$$

we can express it in local coordinates $D_k\mathcal{R}_{ij} = D_i\mathcal{R}_{kj}$. In this case, $Ric(g)$ is a Codazzi tensor . In particular, any Riemannian manifold with Ricci parallel tensor $Dr = 0$ (i.e. $D_i\mathcal{R}_{kj} = 0$) has an harmonic curvature. Indeed, the Levi-Civita connexion D is a Yang-Mills connexion on the tangent bundle of M . In this way, the connexion D is a critical point of the Yang-Mills functional

$$\mathcal{YM}(\nabla) = \frac{1}{2}\int_M ||R^\nabla|| dv,$$

where R^∇ is the curvature associated to the connexion ∇ . Notice that the Riemannian curvature must be harmonic for all Einstein manifolds and for all conformally flat manifolds with constant scalar curvature; this result can be deduced from the orthogonal decomposition of the curvature tensor. Moreover, the condition of the curvature harmonicity is, in a way , a generalisation of the Einstein condition of the metric ($Ric(g)$ is a constant multiple of g : $Ric(g) - \frac{R}{n}g = 0$). In particular, this fact shows that every Einstein metric must have a parallel Ricci tensor $Dr = 0$. In general, the latter property fails for a conformally flat metric with constant scalar curvature, as we shall see below. However, in the 3-dimensional case, we get an identity between harmonic curvature metrics and conformally flat metrics with constant scalar curvature.

Bourguignon proved the converse for the compact manifold of dimension 4 and non vanishing signature. But the signature assumption is necessary, as has been shown by Derdzinski [De]. Moreover, Bourguignon has asked about the existence of compact metric with harmonic curvature and non parallel Ricci curvature, as it was wellknown in the non compact case.

Derdzinski has given examples of metrics with harmonic curvature but non parallel Ricci tensor $Dr \neq 0$, [De]. The corresponding manifolds are bundles with fibres N over the circle S^1 (parametrized by arc length t) with allowed warped metrics $dt^2 + h^{4/n}(t)g_0$ on the product $S^1 \times N$. Here, (N, g_0) is an Einstein manifold of dimension $n \geq 3$, and the function

$h(t)$ on the prime factor is a periodic solution of the ODE, established by Derdzinski

$$h'' - \frac{nR}{4(n-1)} h^{1-4/n} = -\frac{n}{4} Ch \qquad \text{for some constant} \qquad C > 0. \tag{12}$$

This function must be non constant, otherwise the corresponding metric has a parallel Ricci tensor.
Notice that the manifolds $S^1 \times N$ are not conformally flat, unless (N, g_0) has constant sectional curvature. Berger has shown that, under the hypothesis, of non negative sectional curvature on a compact Riemannian manifold with harmonic curvature, the Ricci tensor is parallel .
J. Lafontaine proved that the connected sum $Y = \# X_i$ have the same properties, where X_i are manifolds with negative constant scalar curvature. We may find many other examples of such manifolds in the chapter 16 of A. Besse [B].

4.2 Curvature property of the pseudo-cylindric metrics

More generally, Derdzinski established a classification of the compact n-dimensional Riemannian manifolds (M_n, g), $n \geq 3$, with harmonic curvature. If the Ricci tensor $Ric(g)$ is not parallel and has less than three distinct eigenvalues at each point, then (M, g) is covered isometrically by a manifold

$$(S^1(T) \times N, dt^2 + h^{4/n}(t) g_0),$$

where the non constant positive periodic solutions h verify the equation (11). Here (N, g_0) is a (n-1)- dimensional Einstein manifold with positive (constant) scalar curvature.
We get the following result, which gives another metric sharing this curvature property. See [Ch2].

Theorem (4-3)
Consider the product manifold $(S^1(T) \times S^{n-1}, g_0)$ *. Under the condition*

$$T(c) \geq T_1 = \frac{2\pi}{\sqrt{n-2}},$$

on the circle length, the Riemannian curvatures of the associate pseudo-cylindric metrics $g_c = u_c^{\frac{4}{n-2}} g_0$ *are harmonic and their Ricci tensors are non parallel.*
Moreover, any pseudo-cylindric metric may be identified to a Derdzinski metric up to a conformal transformation.

Return now to the Derdzinski metrics descrived above. We remark that, in the locally conformally flat case of this product, the factor N may be identified with the standard sphere S^{n-1} . Hence, these metrics must be conformally flat and will be conformal to the cylindric metric, as S.T. Yau [Y] has shown

Lemma (4-4)
Any warped metric $dt^2 + f^2(t)g_0$ *on the product Riemannian manifold* $(S^1(T) \times S^{n-1}, dt^2 + g_0)$ *must be conformally flat and conformal to a Riemannian metric product* $d\theta^2 + d\xi^2$ *where* θ *is a* S^1*-parametrisation with length* $\int_{S^1} \frac{dt}{f(t)}$.

Thus, any Derdzinski metric may be identified with a pseudo-cylindric metric up to conformal diffeomorphism, let F . Then the metrics are relied

$$dt^2 + f^2(t)d\xi^2 = F^*({u^j}_c^{\frac{4}{n-2}}(dt^2 + d\xi^2)),$$

where ${u^j}_c$ are the pseudo-cylindric solutions belonging to the (same) conformal class.
Indeed, for the metric $dt^2 + f^2(t)d\xi^2$ we can write

$$dt^2 + f^2(t)d\xi^2 = f^2(t)[(\frac{dt}{f})^2 + d\xi^2].$$

After a change of variables and by using the conformal flatness of the product metric, we get

$$dt^2 + f^2(t)g_0 = \phi^2(\theta)[d\theta^2 + d\xi^2]$$

which is conformally flat.
This result is not surprising, because any manifold carying a warped metric product $(S^1 \times N, dt^2 + h^{4/n}(t)g_0)$ with harmonic curvature is not conformally flat unless (N, g_0) is a space of constant curvature. This manifold must be locally conformally equivalent to the trivial product $S^1 \times N$.
This product will be conformally flat only if N has constant sectional curvature.

4.3 Remark concerning the parallelism of the Ricci tensor

Parallelism property of the Ricci tensor have an interest particularly for the conformally flatness case. Indeed, consider any Riemannian manifold (M, g) with a parallel Ricci tensor. This implies in particular, that its Weyl tensor is harmonic: $\delta W = 0$ (in local coordonates $D_k W^h_{ijk} = 0$). For a metric $\tilde{g}$ in the conformal class and with harmonic Weyl tensor $[g]: \ \tilde{g} = e^{2\rho} g$, we get the following equality

$$\tilde{\delta}\tilde{W} = \delta W - \frac{1}{2}(n-3)W(\nabla\rho, ., ., .).$$

Since, the corresponding Ricci tensor is parallel, we then obtain in local coordonates

$$(n-3)W^l_{ijk} D_l \rho = 0.$$

Thus, when (M, g) is not conformally flat, then $\rho = 0$ necessarily (see "The Ricci calculus" of Schouten).

5 On the existence of Derdzinski metrics

5.1 Analysis of the Derdzinski equation

The above identification of the two family of metrics must reveal a similarity between the equations (8) and (12), since they have been the starting point of our study. Indeed, by considering another time the ODE point of view, we are able to analyse all the solutions of equation (12), like as we have made in the preceding section 2. Recall that they correspond to metrics with $Dr \neq 0$.

So, this equation may have a non constant, positive periodic solution. To that end, we make a change of variables $h(t) = \alpha f(\beta t)$, where

$$\alpha = (\frac{(n-1)C}{nR})^{-n/4} \qquad and \qquad \beta = \sqrt{\frac{nC}{4}}.$$

Note that the constant C must be positive to ensure that the equation (12) to have a periodic non constant solution. This change gives the equation

$$f'' - f^{1-4/n} + f = 0. \tag{13}$$

We can verify that this equation satisfies the lemmas (2-6) and (2-7) given below in the last section. They will be used to complete the analysis of the equation (8).
Indeed, we get the functions
$g(f) = f - f^{1-\frac{4}{n}} \quad ; \quad g'(f) = 1 - (1-\frac{4}{n})f^{-\frac{4}{n}}$
and $g''(f) = \frac{4}{n}(1-\frac{4}{n})f^{-1-\frac{4}{n}}$.
Notice that the hypothesis $g''(f) \geq 0$ is only satisfied in dimension $n \geq 4$.
Then, we calculate

$$\Delta(f) = (f-1)[g''(1)g'(f) - g'(1)g''(f)].$$

We get

$$\Delta(f) = (f-1)\frac{4}{n}(1-\frac{4}{n})[1 - f^{\frac{4}{n}} + \frac{4}{n}(f^{-\frac{4}{n}} - f^{-1-\frac{4}{n}})].$$

It follows

$$\Delta(f) = \frac{4}{n}(1-\frac{4}{n})f(1-f^{-1})[1 - f^{\frac{4}{n}} + \frac{4}{n}f^{-\frac{4}{n}}(1-f^{-1})],$$

which is positive.
Hence, we deduce the increase of the period function depending on the energy

of the equation (12). So, this fact allows us to determine the exact number of these Derdzinski metrics. We first remark, that the bifurcation points of the solution family are

$$(T_k, u_k) \quad , \ where \quad u_k = (\frac{(n-1)C}{nR})^{-n/4} \quad and \quad T_k = \frac{2\pi}{\sqrt{C}}.$$

5.2 Existence conditions of the Derdzinski metrics

It appears from the above analysis that, a non constant, periodic solution of the equation (12) exists only if the circle length satisfies the following condition

$$T \geq \frac{2\pi}{\sqrt{C}}. \tag{14}$$

Notice that, we get an infinity of solutions . All are obtained by rotation-translation of the variable.
Moreover, in the case where T satisfies the double inequality

$$2\pi\frac{(k-1)}{\sqrt{C}} \leq T < 2\pi\frac{k}{\sqrt{C}} \quad k \ is \ an \ integer \ > 1, \tag{15}$$

then the equation (8) may admit k rotationnally invariant distinct solutions. Hence, we have improved the lemma 1 of Derdzinski [De] (see also Chapter 16 of A. Besse [Be])

Theorem (5-3)
Consider the warped product $(S^1(T) \times N, dt^2 + h^{4/n}(t)g_0)$ *. The non constant positive periodic solution* $h(t)$ *satisfies the equation (12) and* (N, g_0) *is a (n-1)-dimensional Einstein manifold with positive (constant) scalar curvature* R *. Then, if the circle length* T *satisfies the following inequalities*

$$2\pi\frac{(k-1)}{\sqrt{C}} \leq T < 2\pi\frac{k}{\sqrt{C}} \ , \qquad where \ k \ \ is \ an \ integer > 1,$$

there exist at least k *rotationnally invariant warped metrics*
$dt^2 + h^{4/n}(t)g_0$ *on the product manifold* $S^1(T) \times N$*.*
Moreover, these metrics have an harmonic curvature. Their Ricci tensor are non parallel only if $T \geq \frac{2\pi}{\sqrt{C}}$ *.*
Conversely, if the warped metric of the type $dt^2 + h^{4/n}(t)g_0$ *on the manifold* $S^1(T) \times N$ *has harmonic curvature, then the function* $h(t)$ *on the circle satisfies an ODE (12).*

The condition $T \geq \frac{2\pi}{\sqrt{C}}$, implies the existence of a non constant solution $h(t)$. Otherwise, we have seen above that a trivial product have a parallel Ricci tensor.

On the other hand, we remark that any warped metric $dt^2 + h^{4/n}(t)g_0$ defined by the lemma (4-4) on the product manifold $(S^1 \times S^{n-1}, dt^2 + d\xi^2)$ is conformal to a Riemannian metric product $d\theta^2 + d\xi^2$. Here θ is a new S^1-parametrisation with length $\displaystyle\int_{S^1} \frac{dt}{h^{2/n}(t)}$.
Furthermore, we have seen in a previous paper [Ch1], there exists a analytic deformation in the conformal class $[g_T]$, of any warped metric $g_T = dt^2 + f^2(t)d\xi^2$ on the manifold $(S^1 \times S^{n-1}, dt^2 + d\xi^2)$, $n = 4$ *or* 6 , and satisfying the length condition

$$T = \int_{S^1} \frac{dt}{f(t)} .$$

Notice that the product metric (under the length condition) belongs to the conformal class $[g_T]$. This analytic family of metrics depends on two parameters $(g_{\alpha,\beta})$. They all have a constant positive scalar curvature, and satisfy the condition: $g_{\alpha,0}$ is the warped metric g_T .
Moreover, by using the same argument of [Ch], we are able to extend the latter result to dimension 3. More precisely, a metric on the euclidean space IR^3 for which the rotation group $SO(3)$ acts by isometries, is in fact a warped product as $g_T = dt^2 + f^2(t)d\xi^2$, where $d\xi^2$ is the standard metric on the sphere S^2 . We can verify that its scalar curvature is

$$R = \frac{2 - 2f'^2 - 4ff''}{f^2} .$$

This fact and the connection with the dimension n will be clarified below, in the next section.

Moreover, we know that every conformally flat manifold (M, g) admits a Codazzi tensor which is not a constant multiple of the metric. Let b a such symmetric 2-tensor field; suppose b has exactly 2 distinct eigenvalues λ , μ , has constant trace $tr_g(b) = c$ and is non parallel. Following [De], these conditions give locally

$$M = I \times N \quad \text{with allowed metric} \quad g = dt^2 + e^{2\psi} g_N \qquad (16)$$

and

$$b = \lambda dt^2 + \mu e^{2\psi} g_N, \quad \lambda = \frac{c}{n} + (1-n)ce^{-n\psi}, \quad \mu = \frac{c}{n} + ce^{-n\psi} . \qquad (17)$$

Conversely, for any such data and for an arbitrary function ψ on I (16) defines a Riemannian manifold (M,g) with Codazzi tensor b of this type. If we assume that the positive function $h = e^{\frac{n}{2}\psi}$ satisfies Equation (12)

$$h'' - \frac{nR}{4(n-1)} h^{1-4/n} = -\frac{n}{4} Ch \qquad \text{for some constant} \qquad C > 0,$$

then the Ricci tensor is precisely a Codazzi tensor and, thus it is non parallel. Therefore, (M,g) is isometrically covered by $(R \times N, dt^2 + h^{4/n}(t) g_N)$ where (N, g_N) is an Einstein manifold with positive scalar curvature.

6 Singular points of the pseudo-cylindric metrics: a global estimate

6.1 A bound for the pseudo-cylindric fonctions

In this part, we examine another point of view coming from the explicit expression of these metrics $g^j{}_c = u^j{}_c{}^{\frac{4}{n-2}} g_0$. We can sometimes solve explicitly the Yamabe equation. In particular, when it is reduced to the equation (8) considered above.

Our approach has not only a calculation interest. However, the direct resolution is very instructive. In particular, it gives the nature of the singularities of the solutions , which seems to depend on the dimension value n.

More precisely, for the dimensional cases $n = 3, 4$ and 6 , we are able [Ch1] to resolve the equation (8) by using the elliptic functions. Notice that an elliptic function is a double periodic meromorphic function, having the same number of poles and zeros in a parallelogramme of periods. For these dimensions, the pseudo-cylindric functions $u^j{}_c$ which have (necessarely) a real period T, turn out to be meromorphic, and have only polar singularities. Moreover, T is precisely the period of the integral curve of Equation (8), which is an elliptic curve (see Whittaker-Watson [W-W]).

We recall that any solution must have a singular point; it becomes infinite on approach of a point $p \in \Lambda$. Furthermore, this solution (which is positive), must have two singular points in 0 *and* ∞ , as has been shown by Cafarelli-Gidas-Spruck [C-G-S]. They have given an estimates near a singular point $p \in \Lambda$, that is for r near 0, u satisfies

$$C^{-1}\, r^{\frac{2-n}{2}} \leq u(r) \leq C\, r^{\frac{2-n}{2}}$$

for some constant C , which depends only on n. In others words the metrics $g^j{}_c = (u^j{}_c)^{\frac{4}{n-2}} g_0$ must be cylindrically bounded near $p \in \Lambda$.

R. Schoen have extended this estimate to the whole manifold $S^n - \Lambda_k$, (Theorem (4-3) of [P]). In the sense, these metrics globally are cylindrically bounded. His proof consider two important remarks. Firstly, by a sequence of dilated coordonates $z_i = \lambda_i y$, the correspondant solutions $v_i(y)$ are uniformly bounded on a large ball in IR^n. Secondly, the Harnack inequality ensures the existence of a positive uniform bound for v_i in any compact set.

By another method, in using properties of some classical properties of elliptic functions, we find again the Schoen result for $k = 2$, and give another estimate concerning the derivative. We obtain the following result, [Ch1].

Theorem (6-1)
Consider the pseudo-cylindric metrics ${g^j}_c = ({u^j}_c)^{\frac{4}{n-2}} g_0$ *on the product Riemannian manifold* $(S^1 \times S^{n-1}, dt^2 + d\xi^2)$. *For the dimensional values* $n = 3, 4$ *and* 6 *only, the corresponding pseudo-cylindric fonction* ${u^j}_c$, *may be expressed in terms of elliptic (meromorphic) functions.*
For these dimensions , we get the estimates

$$r^{\frac{n-2}{2}} u(r) \leq C_n(T) \qquad \textit{and} \qquad r^{\frac{n}{2}} \frac{du(r)}{dr} \leq C'_n(T),$$

where the constants C *and* C' *depend on the period* T.
For the other dimensions, the corresponding pseudo-cylindric fonction ${u^j}_c$, *may be expressed in terms of automorphic functions.*

6.2 General remarks

(i) Consider a pseudo-cylindric metric ${g^j}_c = ({u^j}_c)^{\frac{4}{n-2}} g_0$ with $j = 1, 2, .., k-1$. By changing the variable t into a new variable $r(t)$, defined by

$$\frac{dr}{dt} = u^{\frac{2}{n-2}} \quad \textit{and} \quad r(0) = 0 \ ,$$

Mazzeo-Pollack-Uhlenbeck [M-P-U] have given a more simple expression of the period. Their method, which is purely geometric, allows a quite explicit solution of the equation (8). Notice that this new variable represents geodesic distance along the ∇t integral curves. The function $u(r)$ is still periodic, and the period $R(c)$ represents the length of the geodesic. For the dimension $n = 4$, the solution can easily be expressed as inverse sinus function of the elliptic Jacobi function considered above. Then, the new period becomes $R(c) \equiv \pi$.

(ii) Now consider the linear differential equation

$$g\Delta f = Hess(f) - fRic(g) \qquad (E)$$

where $Hess(f)$ denotes the second covariant derivative of f , $\Delta f = trace[Hess(f)]$ and $Ric(g)$ is the Ricci curvature tensor. Since this equation is linear, $f = 0$ is the trivial solution. This equation was originally derived from the linearization of the scalar curvature equation. The scalar curvature is considered as a non-linear map $\mathcal{R}$ of the space of C^∞ riemannian metrics whose derivates are L_2-integrable to the space of C^∞ functions on the manifold M . This map is differentiable and has derivative

$$d\mathcal{R}_g(\phi) = \frac{d}{dt}|_{t=0} \mathcal{R}(g + t\phi) = -\Delta tr\phi + divdiv\phi - \phi.Ric$$

and the adjoint operator $d\mathcal{R}_g^*$ of $d\mathcal{R}_g$ with respect to the inner product is

$$d\mathcal{R}_g^*(f) = -g\Delta f + Hessf - fRic(g).$$

Thus $d\mathcal{R}_g^*$ is not surjective when $d\mathcal{R}_g^*$ has a non-trivial kernel.

The equation (E) may have another (non-trivial) solution. Let (M, g) be the product manifold $(S^1 \times S^{n-1}, dt^2 + d\xi^2)$ where $(S^{n-1}, d\xi^2)$ is the standard (n-1)-dimensional sphere with scalar curvature $\mathcal{R}_0$ and (S^1, dt^2) is the circle of length T parametrized by t. Under the additional geometric condition

$$T = 2\pi\sqrt{\frac{n-1}{R_0}} \ , \tag{18}$$

the equation have a non trivial solution on (M, g) .
This condition corresponds precisely to the degenerate case of the product metric, in the sense that the Yamabe solution is a degenerate critical point of the functional $J(u)$. Moreover, this example (with other non trivial ones) was used to give a counter-example to the Fisher-Marsden conjecture.
Notice that this manifold is conformally flat and has a parallel Ricci tensor. Proposition 6.5 in [M-P-U] asserts that, among the pseudo-cylindric metrics, only the cylindrical (trivial product) one gives a non trivial solution to Equation (E)

$$(E) \qquad g\Delta f = Hess(f) - fRic(g)$$

but under the additional length condition $T = T_k = \frac{2\pi k}{\sqrt{n-2}}$.

7 Curvature of the asymptotic pseudo-cylindric metrics

An interesting and natural question is to ask if any complete Yamabe metric on $S^n - \Lambda_k$ where $\Lambda_k = \{p_1, p_2, ..., p_k\}$ $k > 2$ shares the same curvature properties of the pseudo-cylindric metrics. This problem is not easy, since we do not have enough indications on these metrics, except their asymptotic behaviour on approaching a point $p_l \in \Lambda_k$.

7.1 Characterization of metrics with k singularities

To be more clear, we use a result of [C-G-S] concerning the estimate of positive solutions $u(t, \xi)$, on the cylinder of the following equation

$$4\frac{n-1}{n-2}\Delta_{g_0}u + R(g_0)u - R(g)u^{\frac{n+2}{n-2}} = 0, \tag{19}$$

$$g = u^{\frac{4}{n-2}}g_0 \quad \textit{complete on} \quad S^n - \Lambda_k \quad \textit{and} \quad R(g) = \textit{constant} > 0 \ .$$

By using a reflection argument, they actually assert that [C-G-S], any solution $u(x)$ of equation (19), with a singularity at p , is asymptotic to one of the pseudo-cylindric functions near the point p . Following [P], we deduce that any such solution $u(x)$ corresponds to a solution $u(t, \xi)$ of equation (8) on a domain of $I\!R \times S^{n-1}$.

Then, the corresponding pseudo-cylindric metric is unique.

Let $\bar{g} = [u(t, \xi)]^{\frac{4}{n-2}}g_0$ be a Yamabe metric conformal to the standard metric and ${u^{j,l}}_c(t)$ be the pseudo-cylindric solution at a point $p_l \in \Lambda_k$. The index j is corresponding to one solution having period T. Then, we obtain the following as a corollary of a result of [C-G-S]

Lemma (7-2)
As $t \to \infty$ *we get the following estimate of the Yamabe solution on* $S^n - \Lambda_k$ *near the point* p_l

$$u(t, \xi) = {u^{j,l}}_c(t) + {u^{j,l}}_c(t)\mathcal{O}(e^{-\beta t}) \ ,$$

where ${u^{j,l}}_c(t)$ *is the corresponding pseudo-cylindric solution and* β *is a positive constant*

In the other words, this solution $u(t,\xi)$, which is asymptotic to a pseudo-cylindric function $u^j{}_c(t)$, around a singular point p_l, can be written as $t \to \infty$

$$u(t,\xi) = u^j{}_c(t) + e^{-\alpha t} v(t,\xi), \tag{20}$$

where α is a positive constant, only depending on the cylindric bound and $v(t,\xi)$ is bounded. This fact was originally pointed out by [C-G-S]. But [P], [M-P-U] have made it more clear, by giving a precise estimate of the decay rate α. It may be calculated in terms of spectral data of the operator $\Delta_\epsilon + n$.

Indeed, approximate solutions for (19) which are singular at the points $p_l \in \Lambda_k$ may be constructed by superposing translated and dilated of the singular radial solutions. The k free parameters in the exact solution correspond to the k different dilational parameters. This fact has been precisely generalized by [M-Pa] in a more general context, to approximate solution not only with isolated singularities, but also when Λ is a k-dimensional submanifold $(0 < k \leq \frac{n-2}{2})$.

So, the corresponding metrics on $S^n - \Lambda_k$ are complete and asymptotic at each singular point to a (unique) pseudo-cylindric metric. Moreover, these "asymptotic metrics" are uniquely determined by the Dilational Pohozaev invariant $\mathcal{D}(g,p)$ on any isolated singular point. [P] has calculated its exact value

$$\mathcal{D}(g,p) = 4\frac{n-1}{n-2}\,\omega_{n-1}\,[E(b) + \frac{1}{n}], \tag{21}$$

where $E(b)$ is the Hamiltonian energy etablished in (2-6). Thus, the Dilational Pohozaev invariant $\mathcal{D}(g,p)$ determines the Hamiltonian energies of the model of the translated pseudo-cylindric solutions for the metric g at each point p_l.

7.2 Ricci tensor of the asymptotic pseudo-cylindric metrics

We get the following result

Theorem (7-3)
Any metric $\bar{g}$ of positive constant scalar curvature which is conformal to the standard metric g_0 on the manifold $S^n - \Lambda_k$, where $\Lambda_k = \{p_1, p_2, ..., p_k\}$ $k > 2$, has a non parallel Ricci tensor, except for the standard one.

Remark (7-4)

Near a singular point $p_l,\ l=1,2,...,k$, any Yamabe solutions on the manifold $S^n-\Lambda_k$ have another property related on the period T_l of the corresponding pseudo-cylindric solutions. Consider its asymptotic expression, as $t\to\infty$,

$$u(t,\xi)={u^{j,l}}_c(t)+e^{-\alpha t}v(t,\xi).$$

Let T_l the (minimal) period of the corresponding pseudo-cylindric function ${u^{j,l}}_c(t)$. This function verify the equality

$$u(t+T_l,\xi)-u(t,\xi)=e^{-\alpha t}[e^{-\alpha T_l}v(t+T_l,\xi)-v(t,\xi)].$$

Since $v(t,\xi)$ is bounded, then for any $t>N$ we have

$$|u(t+T_l,\xi)-u(t,\xi)|<\epsilon.$$

8 Proofs

8.1 Proof of Theorem (2-4)

We first remark there is a conformal diffeomorphism between the manifolds $IR^n - \{0\}$ and $IR \times S^{n-1}$ given by sending the point x to $(log|x|, \frac{x}{|x|})$. By using the stereographic projection, we see easily that the manifold $IR \times S^{n-1}$ (which is the universal covering space of $S^1 \times S^{n-1}$) is conformally equivalent to $S^n - \{0, \infty\}$.
Notice that the manifold $S^n - (p, -p)$ with the standard induced metric, can be considered as the warped product

$$]0, \pi[\times S^{n-1} , \quad \text{with allowed metric} \quad dt^2 + \sin^2 t d\xi^2$$

It has been shown by Cafarelli-Gidas-Spruck [C-G-S] (using an Alexandrov reflection argument), that any solution of (2-1) is in fact a spherically symmetric radial function (depending on geodesic distance from either p or $-p$). Any solution of Equation (6) which gives a complete metric on the cylinder $\Re \times S^{n-1}$ is of the form $u(t, \xi) = u(t)$, where $t \in IR$ and $\xi \in S^{n-1}$.
The background metric on the cylinder is the product $g_0 = dt^2 + d\xi^2$. For the convenience, we assume the sphere radius equal to 1.
Therefore, the partial differential equation is reduced to an ODE.
The cylinder has scalar curvature $R(g_0) = (n-1)(n-2)$ and $R(u^{\frac{4}{n-2}} g_0) = n(n-1)$. Thus $u = u(t)$ satisfies

$$\frac{d^2}{dt^2}u - \frac{(n-2)^2}{4}u + \frac{n(n-2)}{4}u^{\frac{n+2}{n-2}} = 0. \tag{22}$$

It follows that a pseudo-cylindric metric (constant scalar curvature metric) on the product $(S^1(T) \times S^{n-1}, g_0)$ corresponds to a T-periodic positive solution of (8) and conversely. The analysis of this equation shows us, that it has only one center $(\alpha, 0)$. This corresponding to the (trivial) constant solution

$$\alpha = (\frac{n-2}{n})^{\frac{n-2}{4}},$$

and all periodic orbits $\gamma_b(t)$ are surrounded by the homoclinic orbit γ_{b_0} . This orbit is parametrized by

$$(u_0(t), v_0(t)) \quad \text{where} \quad u_0(t) = (cosht)^{-\frac{n-2}{2}}.$$

We denote the coordinates of $\gamma_b(t)$ by $(u_b(t), v_b(t)$. When the value b satisfies the condition $1 < b < b_0$, the correspondant orbit is periodic (b_0 corresponds to a periodic solution of null energy, as we shall see below the

definition) .
One may easily remark that two positive T-periodic solutions of (8) having the same energy are translated, and thus give rise to equivalent metrics on $(S^1(T)\times S^{n-1}, g_0)$.
Note that the metric corresponding to the conformal factor u_0 : $g = {u_0}^{\frac{4}{n-2}} g_0$ is incomplete. Then, it is not a pseudo-cylindric metric; for this reason the constant b cannot attain the critical value b_0 .
To prove Proposition (2-4), we need the following result

Lemma (2-5)
The solution family $(T, u_T(t))$ of the ODE (2-4), (where T is the minimal period) has bifurcation points on the values $(T, u_T(t))$ where $T_k = \frac{2\pi k}{\sqrt{n-2}}$ and $u_{T_k} \equiv \alpha$. In this family, there is a curve of non trivial solutions which bifurcates to the right of the trivial one.

This lemma is a classical result of global bifurcation theory (Crandall-Rabinowitz [C-R]). Indeed, let us consider a positive T-periodic solution: if $T \neq T_k$, then the linearized associate equation is non-singular.
We may also deduce from bifurcation theorem, applied to the simple eigenvalues problem, that there is an unique curve of non trivial solutions near the point (T_k, α). In fact, this uniqueness is global. The trivial curve is $u_T \equiv \alpha$.
Moreover, $(\frac{du}{dT})_{T=T_k}$ is an eigenvalue of the linearized associate equation. According to the global bifurcation theory, we assert that the non trivial curves turn off on the right of the singular solution $(T_k, u_{T_k}(t))$. Consequently, when T varies, two non trivial curves never cross.

On the other hand, we find it more convenient to analyze the equation (8), by making a change of variables $u(t) = \alpha w(\beta t)$, where

$$\alpha = (\frac{n-2}{n})^{\frac{n-2}{4}} \quad \text{and} \quad \beta = \frac{n-2}{2}.$$

This change transforms (21) into the equation

$$\frac{d^2}{dt^2} w - w + w^{\frac{n+2}{n-2}} = w'' + g(w) = 0. \tag{23}$$

If $w(t)$ is a solution of (22), then the Hamiltonian energy

$$E(t) = \frac{w'^2(t)}{2} - \frac{w^2(t)}{2} + (n-2)\frac{w^{\frac{2n}{n-2}}(t)}{2n} = \frac{w'^2}{2} + G(w) - \frac{1}{n}$$

is independent on t . Note that the constant solution $w(t) = 1$ has energy $E_1 = -\frac{1}{n}$, which is the smallest value that E can take. Denote by w_b the solution of (22) which has $w(0) = b > 1$ and $w'(0) = 0$. Then,

$$E(b) = -\frac{b^2}{2} + (n-2)\frac{b^{\frac{2n}{n-2}}}{2n} = c - \frac{1}{n}$$

is the energy of this solution . If

$1 < b < b_0$ *where* $b_0 = \frac{1}{\alpha}$ satisfies $E(b_0) = 0$, then $w_b(t)$ is a positive, periodic function of t , oscillating around the value 1 , we have

$$G(w) = \int_0^w g(u)du \geq 0 \quad \text{satisfying} \quad G(a) = G(b) = c,$$

where a belongs to the interval $[0, 1]$. Hence, we get the expression of the period

$$T(c) = \sqrt{2}\int_a^b \frac{dw}{\sqrt{c - G(w)}}$$

Then, to complete the proof of Proposition (2-4), we also need the following result of Chow and Wang.

Lemma (2-6)

The minimal period $T(c)$ *of a (periodic, positive) solution* u_c *of the equation (22) is a monotone increasing function of its energy, when* $c \in [0, \frac{1}{n}]$

Chow-Wang [C-Wa] also have calculated the derivative of the period function, and have found the expression of the derivative

$$T'(c) = \frac{1}{c}\int_a^b \frac{g^2(w) - 2G(w)g'(w)}{g^2(w)\sqrt{c - G(w)}}dw.$$

To prove the preceding Lemma, we need their following result (see corollary (2-5) in [C-Wa])

Lemma (2-7)

Consider the hypothesis on the smooth function g $g(1) = 0$, $g'(1) < 0$, *we suppose in addition that*

$$H(x) = g^2(x) - 2G(x)g'(x) + \frac{g''(1)}{3g'^2(1)}g^3(x) > 0,$$

for all $x \in [0, b_0]$, . *Then,* $\qquad T'(c) \geq 0 \qquad$ *for all* $c \in [0, \frac{1}{n}]$.

Moreover, they show that under the conditions,

$$g''(x) \geq 0 \quad \textit{and} \quad \Delta(x) = (x-1)\Big[g'(x)g''(1) - g'(1)g''(x)\Big] \geq 0$$

the function $H(x)$ satisfies the previous Lemma (2-7) (that is the corollary (3-1) in [C-Wa]).

Now, we apply these results to the function

$$g(x) = -x + x^{\frac{n+2}{n-2}}.$$

So we get obviously

$$g'(1) = \frac{4}{n-2} \quad , \qquad g''(1) = \frac{4(n+2)}{(n-2)^2},$$

then we calculate

$$\Delta(x) = (x-1)\Big[(\frac{(n+2)}{(n-2)}x^{\frac{4}{n-2}} - 1) - \frac{4(n+2)}{(n-2)^2}(\frac{4}{n-2}x^{\frac{6-n}{n-2}})\Big],$$

and we find

$$\Delta(x) = \frac{4(n+2)}{(n-2)^2}(x-1)\Big[(x^{\frac{4}{n-2}} - 1) + \frac{4}{n-2}x^{\frac{6-n}{n-2}}(x-1)\Big],$$

which is positive.

8.2 Proof of Theorem (4-3)

To prove this Theorem, we consider a local chart in this product manifold, where the local coordonates in this chart are $(x^0, x^1, ..., x^{n-1})$, and $x^0 = t$ is the coordinate corresponding to the circle factor S^1. We can write any metric tensor in the conformal class $[g_0]$: $\bar{g} = u^{\frac{4}{n-2}}g_0$, where the C^∞ function u is defined on the circle. Then we obtain the expressions in the local coordonates system

$$\bar{g}_{00} = u^{\frac{4}{n-2}}, \quad \bar{g}_{0i} = 0, \quad \bar{g}_{ij} = u^{\frac{4}{n-2}}(g_0)_{ij}, \tag{24}$$

where $i, j = 1, 2, ..., n-1$.
For the following, we refer to Kobayashi-Nomizu [K-N].

We get the Christoffel symbols $\bar\Gamma^i_{jk}$ associated to the metric $\bar g$. They are defined by the expression

$$\bar\Gamma^i_{jk} = \frac{1}{2}\bar g^{il}\frac{\partial \bar g_{jl}}{\partial x^i} + \frac{\partial \bar g_{il}}{\partial x^j} - \frac{\partial \bar g_{jk}}{\partial x^l}).$$

Then, we obtain

$$\bar\Gamma^0_{jk} = -\frac{2}{n-2}\frac{u'}{u}\bar g_{jk}, \quad \bar\Gamma^i_{j0} = \frac{2}{n-2}\frac{u'}{u}\bar\delta^i_j, \quad \bar\Gamma^0_{00} = \frac{2}{n-2}\frac{u'}{u} \ \textit{and}\ \bar\Gamma^i_{00} = 0$$

The Ricci tensor components are

$$\bar{\mathcal{R}}_{ij} = \mathcal{R}_{ij} - \frac{2\nabla_{ij}u}{u} + \frac{2n}{n-2}\frac{\nabla_i u \nabla_j u}{u^2} - \frac{2}{n-2}\frac{\mid \nabla u \mid^2 + u\Delta u}{u^2} g_{ij}.$$

We have

$$\bar{\mathcal{R}}_{0i} = 0, \qquad \bar{\mathcal{R}}_{00} = 2\frac{n-1}{n-2}[\frac{u''}{u} + \frac{u'^2}{u^2}].$$

The associated scalar curvatures are

$$R(g_0) = (n-1)(n-2) \qquad and \qquad \bar R(u^{\frac{4}{n-2}}g_0) = n(n-1).$$

So, the trace of the Ricci tensor gives the Yamabe equation

$$\bar g^{ij}\bar{\mathcal{R}}_{ij} = \bar{\mathcal{R}} = u^{\frac{-4}{n-2}}(R + \frac{4(n-1)}{(n-2)}\frac{\Delta u}{u}),$$

and the function u satisfies the above equation

$$u'' - \frac{(n-2)^2}{4}u + \frac{n(n-2)}{4}u^{\frac{n+2}{n-2}} = 0.$$

The covariant derivative of the Ricci tensor gives

$$D_k\bar{\mathcal{R}}_{ij} = \frac{\partial \bar{\mathcal{R}}_{ij}}{\partial x^k} - \bar\Gamma^l_{ik}\bar{\mathcal{R}}_{lj} - \bar\Gamma^l_{jk}\bar{\mathcal{R}}_{li},$$

where D denotes the riemannian connexion associated to the metric g. In particular we get

$$D_0\bar{\mathcal{R}}_{00} = \frac{d\bar{\mathcal{R}}_{00}}{dt} - 2\bar\Gamma^k_{00}\bar{\mathcal{R}}_{k0}.$$

Thus, we obtain

$$D_0\bar{\mathcal{R}}_{00} = 2\frac{n-1}{n-2}\frac{u'''}{u} + 2\frac{n+2}{n-2}\frac{u'u''}{u^2} - 4\frac{u'^3}{u^3}.$$

By comparison with (2-4), u should be a periodical function. So, we deduce that, if the function u is non constant, then

$$D_0\bar{\mathcal{R}}_{00} \neq 0.$$

The Ricci curvature of the pseudo-cylindric metrics is not parallel, except for the cylindric one.

8.3 Proof of Theorem (6-1)

Consider the orbits of the autonomous system associated to the equation (8)

$$u'^2 = (\frac{n-2}{2})^2\left[u^2 - u^{\frac{2n}{n-2}}\right] + \bar{c} \tag{25}$$

these orbits are closed only if the energy constant satisfies the following condition

$$-\frac{2}{n}(\frac{n-2}{2})^2(\frac{n-2}{n})^{\frac{n-2}{2}} < \bar{c} < 0. \tag{26}$$

In connection with the energy constant c, used in the proposition (2-4), we obtain the relation

$$\bar{c} = -2\alpha\beta^2 c = -\frac{(n-2)^2}{2}(\frac{n-2}{n})^{\frac{n-2}{2}} c,$$

where the constants are as above $\alpha = (\frac{n-2}{n})^{\frac{n-2}{2}}$ *and* $\beta = \frac{n-2}{2}$ (we recall that the constant c must varying in the interval $[0, \frac{1}{n}]$).
We remark that for the n-values $n = 4$ *or* 6, all the orbits are elliptic curves, which may be parametrized only by 2-periodic meromorphic functions (i.e. elliptic functions).
More exactly, a direct calculation gives the explicit solutions, for $n = 4$

$$u_{\bar{c}}(r) = \frac{1}{\sqrt{2-\bar{c}^2}}\ dn_{\bar{c}}(\frac{r+\beta}{\sqrt{2-\bar{c}^2}}) \tag{27}$$

where $dn_{\bar{c}}$ is the elliptic Jacobi function

$$dn_{\bar{c}}(x) = 1 - \frac{\bar{c}^2}{2}x^2 + \frac{\bar{c}^2(4+\bar{c}^2)}{6}x^4 + ...$$

of real period

$$T(\bar{c}) = 2\sqrt{2-\bar{c}^2}\int_0^{\pi/2}\frac{d\theta}{\sqrt{1-\bar{c}^2\sin^2\theta}}.$$

These Jacobi functions $dn_{\bar{c}}(r)$ satisfy the differential equation

$$(\frac{d}{dr}dn)^2 = (1-dn^2)(dn^2+\bar{c}^2-1)$$

with $dn_{\bar{c}}(0) = 1$, Whittaker-Watson [W-W].

We obtain analogous results for the case $n = 6$, where the solution may be expressed in terms of the Weierstrass function $\wp(r)$. We recall the equation satisfied by the Weierstrass function

$$\wp'^2(r) = 4\wp^3(r) - g_2\wp(r) - g_3,$$

and its expression is

$$\wp(r) = \frac{1}{r^2} + \sum_{0<p} a_p(g_2, g_3) r^p.$$

In particular we deduce that, $r^2\wp(r)$ is an entire function in an elementary parallelogram of periods. Then, it is bounded.
So, we get the following relation

$$u_{\bar{c}}(r) = \frac{1}{3} - \wp(r+\beta; g_2, g_3), \tag{28}$$

where $g_2 = \frac{4}{3}$ and g_3 which depends on the parameter $\bar{c}$: $(g_3 = -\frac{8}{27} - \bar{c})$ satisfies the condition $g_3 > -(\frac{2}{3})^3$.
This latter inequality, corresponding to the (25), insures us that the function $\wp(r)$ has a real period. Furthermore, if $r = 0$ is a singular point, then we can take the value $\beta = 0$.
It is known that this elliptic function has a double pole at the origin and two zeros (in a elementary parallelogram of periods). Thus, its derivative has 3 zeros and a pole of order 3, [W-W].

Moreover, it is known when r increases in the interval $]0, \frac{T}{2}[$, then the Weierstrass function $\wp(r)$ decreases monotonically in $]+\infty, \wp(\frac{T}{2})[$. So, we obtain explicitly for $n = 6$

$$r^2 u(r) \leq T^2 u(\frac{T}{2}).$$

Also, for the dimension $n = 3$ or $n = 4$, we get an analog

$$r^{\frac{n-2}{2}} u(r) \leq T^{\frac{n-2}{2}} u(\frac{T}{2}).$$

Examine now the general case: to study the integral curves of our equation (24) and their algebraic nature; it seems that it will depend on the parity of the dimension. For this, we consider separately two hypotheses concerning the parity.

(i) Suppose that the dimension n takes an odd value $n = 2p + 1$. Using the change of function

$$u = v^{-\frac{n-2}{2}} ,$$

we get $u'^2 = (\frac{n-2}{2})^2 v^{-n} v'^2$ and, after the replacement, Equation (24) becomes

$$(\frac{n-2}{2})^2 v^{-n} v'^2 = (\frac{n-2}{2})^2 [v^{-(n-2)} - v^{-n}] + \bar{c} ,$$

and we obtain the analogous equation for v

$$v'^2 = v^2 - 1 + (\frac{2}{n-2})^2 \bar{c}\ v^n . \tag{29}$$

By referring to the geometry of algebraic curves (Shafarewitch [Sh], for any odd value $n \neq 3$, the integral curve of this equation is known to be represented by a hyperelliptic curves. Whose coordonates are connected by the algebraic relation

$$F(x,y) = y^2 - x^2 + 1 - (\frac{2}{n-2})^2 \bar{c}\ x^n = 0 .$$

Under condition (26) on $\bar{c}$, the integer p is precisely the genus of such curves if the polynomial $P(x) = x^2 - 1 + (\frac{2}{n-2})^2 \bar{c}\ x^n$ is without multiple roots. Otherwise, the genus fails (which corresponds to the extremal values of $\bar{c}$).
Moreover, as we may easily remark, the curve is elliptic only for the value $n = 3$. For this particular case, we get the equation

$$(a) \qquad v'^2 = v^2 - 1 + 4\bar{c}\ v^3.$$

This equation has a elliptic functions as solutions. More exactly, under the condition (26) on $\bar{c}$,

$$-\frac{1}{6\sqrt{3}} < \bar{c} < 0$$

we get the solution

$$v_{\bar{c}}(r) = \frac{1}{\bar{c}}[\wp(r+\beta; g_2, g_3) - \frac{1}{12}],$$

where $\wp$ is the Weierstrass function relative to $g_2 = \frac{1}{12}\ and\ g_3 = \bar{c}^2 - \frac{1}{216}$. This function has two simples zeros in an elementary parallelogram of periods, one of them is real. We choice β such that $wp(\beta) = \frac{1}{12}$. So, we get $v_{\bar{c}}(0) = 0$. We deduce that the function $u_{\bar{c}} = v_{\bar{c}}^{-\frac{1}{2}}$ has a pole of order $\frac{1}{2}$.
Otherwise, for any odd value of the dimension $n > 3$, the corresponding integral curve of (29) does not admit a parametric representation in terms of

meromorphic functions, because it has genus $p > 1$.
Equivalently, the algebraic relation $F(x,y) = 0$ cannot be uniformized by meromorphic functions in the finite plane. In fact, this curve may be uniformized

$$x = A(r) \quad ; \quad y = B(r),$$

where A and B are automorphic functions which are meromorphic but have only a restricted domain of existence with a natural boundary.

(ii) - Now, consider the case where the dimension n takes an even value. This seems to be harder than the latter case; in particular determining the genus of the integral curve is less easy. We use a more appropriate change

$$u = v^{-\frac{n-2}{4}} ,$$

we get

$$u'^2 = (\frac{n-2}{4})^2 v^{-\frac{n+2}{2}} v'^2$$

and the equation (24) becomes

$$(\frac{n-2}{4})^2 v^{-\frac{n+2}{2}} v'^2 = (\frac{n-2}{2})^2 [v^{-\frac{n-2}{2}} - v^{-\frac{n}{2}}] + \bar{c},$$

then we deduce the analogous equation for v

$$v'^2 = 4v^2 - 4v + (\frac{4}{n-2})^2 \bar{c}\, v^{\frac{n+2}{2}}. \tag{30}$$

To know the algebraic nature of the integral curve of this equation, we proceed as before. Remark that, for the values $n = 4$ *or* 6 , the associated integral curve for this equation is an elliptic one (here the genus is $p = 1$) . However, these two cases have been treated before.
Otherwise, for any even value $n \geq 8$, the integral curve of (29) is a hyperelliptic one, if its second member is without multiple roots.
The genus p of this curve depends on the parity of $\frac{n}{2}$; it is $\left[\frac{n}{4}\right]$, the integer part of $\frac{n}{4}$.
It was proved in [Sh], that the case where the polynomial P has even degree, reduces to that of odd degree.
Moreover, this transformation preserves the genus. For this reason the curves corresponding to $n = 4$ *and* 6 have genus $p = 1$, likewise for those corresponding to $n = 8$ *and* 10 , which have genus $p = 2$, and so on.
Generally speaking, every integral curve of (28) and (29) admits $2p$ abelian periods. One of them is real, and this one is interesting for our problem.

On the other hand, except for the cases $n = 3, 4$ and 6 , the general solution of the equations (29) and (30) may be expressed in terms of automorphic functions. These functions may be regarded as a parametric representation of the corresponding integral curves.
It is known that this solution is an automorphic function of a Fuchsian group of the first kind, all of whose homographic substitutions are hyperbolic. The fundamental region (for the same genus p) is contained in the interior of the limit circle.
Notice that every limit point is necessary an essential singularity for any (non degenerate) solution.
In general, it seems to be very difficult to obtain explicit formulas, the theory of elliptic functions is rather exceptional.

8.4 Proof of Theorem (7-3)

To prove this, we need the asymptotic property of these metrics. In particular, the estimate (20) play an essential role.
Notice that, this manifold $(S^n - \Lambda_k, \bar{g})$ is locally conformally flat. The condition of constant scalar curvature implies that its Riemannian curvature is harmonic. We get the converse only for the dimension $n = 3$.

We proceed as in the proof of Theorem (4-3). Consider a local chart (t, ξ) of a point in $S^n - \Lambda_k$, here $t = x^0$ and $\xi = (x^1, x^2, ..., x^{n-1})$.
In this local coordonates system, we obtain the components of the tensor metric

$$\bar{g}_{00} = u^{\frac{4}{n-2}} \quad , \qquad \bar{g}_{0i} = 0 \quad , \qquad \bar{g}_{ij} = u^{\frac{4}{n-2}}(g_0)_{ij}.$$

The Christoffel symbols $\bar{\Gamma}^i_{jk}$ associated to the metric tensor $\bar{g}$ are

$$\bar{\Gamma}^i_{j0} = \frac{2}{n-2}\frac{1}{u}[\frac{\partial u}{\partial x^j} + \frac{\partial u}{\partial x^i}\delta^i_j] \qquad \bar{\Gamma}^0_{00} = \frac{2}{n-2}\frac{1}{u}\frac{\partial u}{\partial t}$$

Then we get components of the Ricci tensor

$$\bar{\mathcal{R}}_{0i} = -\frac{2\nabla_{i0}u}{u} + \frac{2n}{n-2}\frac{1}{u^2}\frac{\partial u}{\partial x_i}\frac{\partial u}{\partial t},$$

$$\frac{\bar{\mathcal{R}}_{00} = 2\frac{n-1}{n-2}[\frac{1}{u}\partial^2 u}{\partial t^2 + \frac{1}{u^2}(\frac{\partial u}{\partial t})^2].}$$

D denotes the Riemannian connexion associated to the metric $\bar{g}$.
Considering now Lemmas (6-2) and (6-3) which imply that

$$\frac{\partial u}{\partial x^i} = e^{-\alpha t}\frac{\partial v}{\partial x^i} \quad and \quad \nabla_{i0}u = -\alpha e^{-\alpha t}\frac{\partial v}{\partial x^i} + e^{-\alpha t}\frac{\partial^2 v}{\partial x^i \partial t}.$$

Recall that $v = v(t, \xi)$ is a bounded function, and α is a positive constant.

Thus, we can write

$$\bar{\mathcal{R}}_{0i} = e^{-\alpha t} f(t, x_1, x_2, ..., x_{n-1})$$

where f is bounded .

Consequently, we may calculate the components of the covariant derivative

$$D_0 \bar{\mathcal{R}}_{00} = 2 \frac{n-1}{n-2} \frac{{u_c}'''}{u_c} + 2 \frac{n+2}{n-2} \frac{{u_c}' u''}{{u_c}^2} - 4 \frac{{u_c}'^3}{{u_c}^3} - e^{-\alpha t} g(t, x_1, x_2, ..., x_{n-1})$$

where g is a bounded function.

Therefore, since t is large, we conclude that

$$D_0 \bar{\mathcal{R}}_{00} \neq 0.$$

REFERENCES

[A1] T.Aubin *Meilleures constantes dans le théorème d'inclusion de Sobolev* J. Of Funct. Anal., 32, 148-174, (1979).

[Be] A.L.Besse *Einstein Manifolds* Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band 10. Springer. New-York, Berlin (1987).

[C-G-S] L.Cafarelli, B.Gidas and J.Spruck *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev Growth* Comm. Pure and Appl. Math., vol XLII (1989), 271-297.

[Ch-W] R.Chouikha and F.Weissler *Monotonicity properties of the period function and the number of constant positive scalar curvature metrics on* $S^1 \times S^{n-1}$ Prepubl. Math., Universite Paris-Nord, n° 94-14, (1994).

[Ch1] R.Chouikha *Famille de métriques conformèment plates et régularité* C.R. Math. Reports. Acad Sc. of Canada, vol XIV (1992) 257-262.

[Ch2] A.R. Chouikha *On the properties of certain metrics with constant scalar curvature* Diff. Geom. and Appl., Sat. Conf. of ICM in Berlin, (1999), Brno, 39-46.

[Ch3] A.R. Chouikha *Nonparallelism of the Ricci tensor of pseudo-cylindric metrics* Geom. Dedicata 93 (2002), 107–112.

[Ch4] A.R. Chouikha *Existence of metrics with harmonic curvature and non parallel Ricci tensor* Balkan J. Geom. Appl. 8 (2003), no. 2, 21–30.

[C-Wa] S.N.Chow and D.Wang *On the monotonicity of the period function of some second order equations* Casopi pro pestovani matematiky, 111 (1986), 14-25.

[C-R] M.Crandall and P.Rabinowitz *Bifurcation from simple eigenvalues* J. Functional Analysis 8, (1971), 321-340.

[De] A.Derdzinski *Classification of certain compact Riemannian manifolds with harmonic curvature and non parallel Ricci tensor* Math. Z., 172, (1980), 277-280.

[De1] A. Derdzinski *On compact Riemannian manifolds with harmonic* *curvature* Math. Ann., vol 259, (1982), 145-152.

[H-V] E.Hebey and M.Vaugon *Meilleures constantes dans le théoème d'inclusion de Sobolev et multiplicités pour les problèmes de Nirenberg et de Yamabe* Indiana Univ. Math. J., 41, (1992), 377-407.

[K-N] S.Kobayashi and K.Nomizu *Foundations of Differential Geometry* vol. 2. New York-London, Interscience (1966).

[L-P] J.M.Lee and T.M.Parker *The Yamabe problem* Bull. Amer. Math. Soc., 17, (1986) p.37-91.

[M-P-U] R.Mazzeo, D.Pollack and N.K.Uhlenbeck *Moduli spaces of singular Yamabe* J. Amer. Math. Soc. 9 (1996), no. 2, 303–344.

[M-Pa] R.Mazzeo and F.Pacard *Constant scalar curvature metrics with isolated singularities* MSRI, dg-ga server, Duke Math. J. 99 (1999), no. 3, 353–418.

[Pa] F.Pacard *The Yamabe problem on subdomains of even dimensional spheres* Topol. Methods Nonlinear Anal. 6 (1995), no. 1, 137–150.

[P] D.Pollack *Compactness results for complete metrics of constant positive scalar curvature on subdomains of S^n* Indiana Univ. Math. J., vol 42, (1993), 1441-1456.

[Sc] R.Schoen *Variational theory for the total scalar curvature functional for Riemannian metrics and related topics* Lectures Notes in Math., 1365, Springer-Verlag (1989), 120-154.

[Sc1] R.Schoen *On the number of constant scalar curvature metrics in a conformal class* Differential Geometry: A symposium in honor of M. Do Carmo. Wiley, (1991), 311-320.

[Sh] I.C.Shafarewitch *Basic algebraic geometry* Springer-Verlag, (1977).

[W-W] E.T.Whittaker and G.N.Watson *A course of modern analysis* Cambridge, (1963).

[Y] S.T.Yau *Remarks on conformal transformations* J. Diff. Geom., 8, (1973), 369-381.